

小心登革熱 齊來把蚊滅

Beware of Dengue Fever



4

花瓶和水養植物須每星期最少換水一次，並清除花瓶、花盆及水桶等器皿內的積水；

Change water for flowers and plants at least once a week. Remove stagnant water in containers like vases, flower pots and buckets;



5


確保貯水容器、貯水箱及水井蓋密；
Tightly cover all water containers, water storage tanks and wells;



6

保持溝渠暢通，避免淤塞；

Keep all drains free from blockage;



7

將凹陷的地面填平；及

Level all defective ground surfaces; and



8



停車場內的防撞車胎須鑽上大孔，以防止積水。

Make large holes in the tyres used as anti-bumping measure in car park to prevent accumulation of water.

查詢 Enquiries

如欲得到有關防治白紋伊蚊的進一步資料，可致電食物環境衛生署**防治蟲鼠事務諮詢組 (3188 2064)**。

For further information on how to control and prevent *Aedes albopictus*, please contact the Pest Control Advisory Section on 3188 2064 of the Food and Environmental Hygiene Department.



中文網站



English website

食物環境衛生署出版 政府物流服務署印
Published by the Food and Environmental Hygiene Department
Printed by the Government Logistics Department (07/2024)





簡介

登革熱是一種由伊蚊傳播的疾病，常見於熱帶和亞熱帶地區。香港夏天炎熱多雨，非常適宜蚊蟲繁殖。蚊蟲主要在雨季開始滋生，並會持續至年底。要預防登革熱的傳播，市民應保持屋邨環境衛生，以防蚊蟲滋生。

Introduction

Dengue fever, an aedine mosquito-borne disease, commonly occurs in tropical and subtropical regions. In Hong Kong, hot summers are accompanied by heavy rainfall, providing favourable breeding grounds for mosquitoes. Mosquito breeding usually begins in the rainy season and continues till the end of the year. To prevent the transmission of dengue fever, the public is advised to keep the estates and the living environment clean against mosquito breeding.

登革熱


登革熱是由過濾性病毒引起的，症狀包括高燒、嚴重頭痛、肌肉及關節痛、眼窩後疼痛及出疹等。潛伏期約為三至十四日。首次感染的病



情一般較輕，但如果隨後再感染其他血清型的登革熱病毒，則有可能出現重症登革熱，嚴重者甚至會死亡。

登革熱的傳播途徑

登革熱病毒是經由胸背有白線和腳部有白間的白紋伊蚊傳播。人被帶有登革熱病毒的白紋伊蚊叮咬後，便可能受感染。市民應時刻保持警覺，防止登革熱的散播。



Dengue Fever

Dengue fever is a viral illness characterised by high fever, severe headache, muscle and joint pain, pain behind the eyes and rash. The incubation period is about 3 to 14 days. Symptoms are usually

mild on the first infection. Subsequent infections with other serotypes of dengue virus can cause severe dengue, and even death.

防治蚊患的方法

要防止蚊蟲滋生，最重要是清除積水。食物環境衛生署建議屋邨管理公司採取以下的措施，以杜絕蚊患：

1

委派員工專責防治蚊患工作：

Appoint designated staff for mosquito prevention and control;

2

最少每星期檢查屋邨範圍一次， 以清除積水：

Inspect the housing estates at least once a week to remove stagnant water;



3

白紋伊蚊喜歡在範圍細小的積水產卵，因此要適當棄置垃圾，如將汽水罐及空飯盒放於有蓋垃圾桶內，以免積水：



As Aedes albopictus may lay eggs in small pools of stagnant water, refuse such as soft drink cans and empty lunch boxes should be properly disposed of in covered litter containers to prevent accumulation of stagnant water;

Control Measures against Mosquitoes

Removal of stagnant water is of prime importance to the prevention of mosquito breeding. The following measures are recommended to the management agencies of housing estates to prevent mosquito breeding: